
# Foot-and-Mouth Disease for Agricultural Shows

**£** What is Food-and-Mouth Disease?

- Highly contagious viral illness affecting cloven-hoofed animals
- Symptoms: Painful blisters, lameness, drooling, reduced appetite and movement
- Economic impact: Trade restrictions, long-term economic losses







- Incubation period: 2–14 days (animals may appear healthy but still be infectious
- Disease spreads quickly via animal movement, contaminated clothing, equipment and vehicles
- Recent outbreaks confirmed in KZN, Gauteng and Mpumalanga. Western Cape remains FMD-free

**Description Legal and Biosecurity Requirements for Agricultural Shows** 

#### **Before the show**

Origin/Owner Declaration Form (indicating origin)

### If you suspect FMD

- Report any suspicion to your local State Veterinarian
- Do not move the animal
- Contact details: www.elsenburg.com

Summary of your responsibilities

#### Show organisers:

- Ensure forms submitted, manage pens, enforce hygiene
  Animal owners:
- Bring healthy animals, provide documentation, isolate animal post-show, avoid risky purchases

#### **Buyers:**

 Agree to quarantine new animals and monitor for symptoms

- and health status)
- Veterinary Certificate (recommended)
- Avoid high-risk animals

#### During the show

- No contact between animals from different farms.
- Avoid shared handling of animals
- Provide footbaths and disinfectant stations
- Disinfect tools, pens and vehicles

#### After the show

- Buyer/Destination Declaration Form
- Mandatory 28-Day quarantine isolation period
- Clean and disinfect all equipment and vehicles

<b>T</b> <sup>++</sup> Disinfectants effective against FMD			
Active Ingredient	<b>Concentration</b> <b>for FMDV (%)</b>	Contact time (minutes)	Important to note
Sodium hydroxide (Caustic soda)	2	10	Add powder to water an not water to powder. Corrosive.
Sodium carbonate (Soda ash)	4	10	Mildly corrosive.
Citric acid	3	Porous surfaces: 30 min. Non-porous surfaces: 15 min	Clean well before applyin
Acetic acid (Household vinegar (4-8% acetic acid))	2	10	Clean well before applyin
Sodium hypochlorite (NaClO) (Household bleach (approx. 3% sodium hypochlorite in South Africa)	0.3	Porous surfaces: 30 min. Non-porous surfaces: 15 min.	Use immediately after mixing the solution. Add NaClO to water. Corrosive.
Potassium peroxymono-sulphate (Virkon S)	1	10	



Scan the QR code or visit:

https://www.elsenburg.com/wp-content/uploads/2025/06/Government-Gazette.pdf



## Expert veterinary services, delivered



FOR YOU